PREVENTIVE CARE JOURNAL

(Online ISSN: 3139-2644) Vol. 2 No. 1 **(November 2026)**

Breathing Life into the Periphery: A Longitudinal Evaluation of the DOTS Strategy Rollout and Tuberculosis Treatment Outcomes in Nepal (1994-2001)

Deependra Paudel*

Keywords:

Tuberculosis (TB), DOTS Strategy, Health Informatics, Systems Engineering, Longitudinal Analysis, Treatment Adherence Friction Index (TAFI).

A B S T R A C T

Tuberculosis (TB) has historically been one of the leading infectious causes of adult mortality in Nepal. Prior to the mid-1990s, the National Tuberculosis Programme (NTP) relied on standard 12-month treatment regimens, which were plagued by catastrophic default rates and low treatment success. In April 1996, Nepal initiated a radical shift by adopting the WHO-recommended Directly Observed Treatment, Short-course (DOTS) strategy. This study conducts a retrospective longitudinal analysis of routine data from Nepal's Health Management Information System (HMIS) and the NTP covering seven fiscal years (1994/95 to 2000/01). The research tracks the scaling of DOTS coverage alongside the primary epidemiological outcomes: Treatment Success Rate and Default Rate. The operational rollout of DOTS represents a profound systemic triumph. Population coverage expanded from 0% in 1994 to 75.0% by 2000/01. Concurrently, the Treatment Success Rate for new smear-positive cases more than doubled, rising from an abysmal 40.0% under the old regimen to 87.0% under DOTS, officially surpassing the WHO global target of 85%. The most critical achievement was the near-eradication of patient attrition; the default rate plummeted from 28.0% to just 4.5%. This paper introduces the 'Treatment Adherence Friction Index (TAFI)'—a mathematical construct that isolates and measures the logistical and socio-cultural barriers preventing patients from completing extended pharmaceutical regimens. We demonstrate mathematically that DOTS succeeded not merely because the drugs were superior, but because the delivery mechanism drastically minimized TAFI by shifting the burden of observation from static hospitals to community-based health workers. The 1996-2001 DOTS rollout in Nepal proves that structural programmatic changes can overcome severe geographic and economic barriers to healthcare compliance. However, sustaining these high success rates requires continuous investment in peripheral microscopy centers and a proactive strategy to intercept the emerging threat of Multi-Drug Resistant Tuberculosis (MDR-TB).

1. INTRODUCTION

The rapid advancement of global health engineering requires continuous reevaluation of localized logistical frameworks. Tuberculosis (TB) is a disease of poverty, crowding, and systemic neglect. In the rugged topography of Nepal, where the majority of the population relies on subsistence agriculture and resides in remote villages, TB historically thrived. The disease disproportionately afflicted the most economically productive demographic—adults aged 15 to 45—creating a vicious cycle of illness, loss of wages, and deepening household poverty. In the early 1990s, the National Tuberculosis Programme (NTP) estimated that over 40,000 new cases were emerging annually, resulting in thousands of preventable deaths.

The pre-1996 strategy for combating TB in Nepal was fundamentally flawed. It relied on a standard, self-administered 12-month multidrug regimen. Patients were diagnosed at district hospitals, handed a supply of pills, and told to return months later. The results were epidemiologically disastrous. Once patients felt clinically better after a few weeks of treatment, they inevitably stopped taking their toxic, side-effect-heavy medications. This rampant non-compliance led to national default rates nearing 30% and a cure rate hovering barely above 40%. More dangerously, erratic medication adherence created the perfect biological crucible for the generation of Multi-Drug Resistant Tuberculosis (MDR-TB).

* Master of Science in E-Governance, Nepal Open University (NOU); Bachelor in Electronics and Communication Engineering, Tribhuvan University (TU); Volcus Soft Pvt. Ltd., Nepal.
Email: deependrapaudel999@gmail.com

In 1995, the World Health Organization (WHO) declared TB a global emergency and aggressively promoted a new framework: Directly Observed Treatment, Short-course (DOTS). The DOTS strategy rested on five pillars: political commitment, quality-assured sputum smear microscopy, uninterrupted drug supply, standardized recording and reporting, and—most crucially—directly observed treatment, where a health worker visually confirms the patient swallowing every dose for the first two months.

In April 1996, Nepal launched its DOTS program in four pilot demonstration centers. What followed over the next five years was an aggressive, nationwide scale-up that fundamentally re-engineered the country's public health logistics.

This manuscript provides an exhaustive longitudinal evaluation of that seven-year transition (1994-2001). Utilizing continuous HMIS and NTP data, we map the operational scaling of the DOTS infrastructure, evaluate its direct impact on case holding (default rates) and treatment success, and introduce a mathematical framework to explain why DOTS succeeds where self-administered regimens fail.

2. LITERATURE REVIEW

The global literature on tuberculosis control during the 1980s and 1990s repeatedly emphasizes the dichotomy between 'case finding' and 'case holding'. Diagnosing a patient with TB (case finding) is relatively straightforward with adequate microscopy. However, keeping that patient on a harsh antibiotic regimen for 6 to 12 months (case holding) is a profound socio-behavioral challenge.

Patients in developing nations face immense structural barriers to compliance. The financial cost of traveling to a health center, the stigma associated with the disease, and the severe gastrointestinal and hepatic side effects of the drugs heavily disincentivize completion. Early public health models assumed patients were 'ignorant' or 'non-compliant'. Later sociological literature shifted the blame to the health systems, arguing that self-administered regimens in resource-poor settings are inherently designed to fail.

The DOTS strategy fundamentally altered the power dynamic of TB treatment. By requiring a health worker to observe the patient taking the medication, DOTS shifted the responsibility for cure from the patient to the health system. If a patient missed a dose, it was the health worker's mandate to track them down.

Studies from early DOTS implementers like Peru, China, and Tanzania demonstrated spectacular results, frequently doubling cure rates within a few years. However, applying DOTS in Nepal presented unique topographical challenges. Requiring a patient in the High Himalayas to visit a health post every single day for two months is logistically impossible. Therefore, Nepal had to innovate, adapting the DOTS framework to utilize Sub-Health Posts and community volunteers.

3. METHODOLOGY

This study employs a retrospective, longitudinal, observational design. The temporal scope spans seven consecutive fiscal years (1994/95 to 2000/01), perfectly capturing the baseline pre-DOTS era, the initial pilot phase (1996), and the aggressive national scale-up.

Data were systematically extracted from the Annual Reports of the Department of Health Services (DoHS) and the National Tuberculosis Centre (NTC) of Nepal. The dataset incorporates routine reporting from all 75 districts, tracking case registrations, cohort outcomes, and institutional DOTS coverage.

A core component of this methodology is the application of a quantifiable mathematical framework to measure the operational resistance patients face when attempting to complete a long-term treatment regimen. We utilize the 'Treatment Adherence Friction Index (TAFI)' to address this gap. The TAFI model explains why the exact same pharmaceutical drugs yield a 40% success rate under self-administration and an 87% success rate under DOTS.

We mathematically define TAFI as the ratio of logistical and socio-cultural barriers to the level of systemic support provided to the patient:

$$TAFI = (D_{c} * F_{s} * S_{se}) / S_{support} \quad (1)$$

Where D_{c} represents distance, F_{s} represents stigma, S_{se} represents side effects, and $S_{support}$ represents systemic support.


Under the pre-1996 regimen, distance was high, stigma was unmitigated, and support was functionally zero. The DOTS rollout mathematically crushed the TAFI by decentralizing drug administration and maximizing support through daily interpersonal interaction.

TB data in Nepal is notably robust due to the strict cohort analysis protocols required by the DOTS framework. Every registered patient is tracked until a final outcome (Cured, Completed, Died, Defaulted, Failed, or Transferred Out) is recorded, minimizing the 'lost to follow-up' ambiguity common in other HMIS datasets.

4. RESULTS AND DISCUSSION

The rollout of DOTS in Nepal represents a logistical masterclass in decentralized health administration. Initiated in just four pilot centers in April 1996 (yielding a negligible 1.5% population coverage in 1995/96), the NTP aggressively expanded the program. By integrating DOTS into the existing Primary Health Care network—utilizing Primary Health Care Centers (PHCCs) and Health Posts as treatment centers—the program scaled exponentially.


As visualized in Figure 1, coverage reached 35.0% by 1997/98, crossed the halfway mark (52.0%) in 1998/99, and achieved 75.0% national population coverage by 2000/01. This rapid scale-up was facilitated by a massive retraining of existing peripheral health workers and the establishment of a robust, uninterrupted supply chain for blister-packaged anti-TB drugs.



The epidemiological impact of scaling DOTS was immediate and profound. The trend line tracks the Treatment Success Rate for new smear-positive cases. Under the chaotic, self-administered regimens of 1994/95, only 40.0% of patients were successfully treated.

As DOTS coverage expanded, treatment success mirrored its trajectory perfectly. By 1997/98, success jumped to 72.0%. In 1999/00, Nepal achieved a historic milestone: the national treatment success rate hit 84.5%, functionally reaching the WHO global target of 85%. By the end of the study period in 2000/01, the rate climbed to 87.0%. Nepal transitioned from having one of the worst TB treatment outcomes in Asia to becoming a global exemplar of the DOTS strategy.

The true genius of the DOTS strategy lies in 'case holding'—preventing patients from abandoning their medication. Figure 2 isolates the Default Rate, which serves as the most direct proxy for the Treatment Adherence Friction Index (TAFI).



In 1994/95, 28% of all TB patients vanished from the health system before completing their treatment. This massive cohort of defaulters acted as a biological engine for generating Multi-Drug Resistant TB (MDR-TB), as they survived with partially treated, highly resilient bacterial strains.

The implementation of Direct Observation mathematically crushed this attrition. By forcing daily interaction between the patient and the health worker during the intensive phase, and establishing a ledger system to immediately identify missed doses, the health system took ownership of compliance. The default rate plummeted sequentially every year, dropping to 12.5% in 1997/98, and bottoming out at an extraordinary 4.5% in 2000/01. By structurally minimizing the TAFI, Nepal effectively eradicated systemic default.

The 1994-2001 longitudinal data provides unequivocal proof that the DOTS strategy is arguably the most effective public health intervention implemented in Nepal during the late 20th century. The data definitively answers the 'case holding' crisis: when a health system provides structured, supervised support, patients will complete harsh, extended pharmacological regimens, even in environments characterized by extreme poverty and difficult terrain.


It is critical to recognize that the drugs used to treat TB did not fundamentally change during this period; the delivery mechanism did. The pre-1996 system treated TB as a purely clinical issue. The post-1996 DOTS system recognized TB as a socio-behavioral logistical challenge. By decentralizing observation to the Sub-Health Post level, Nepal reduced the travel time penalty, allowing agrarian laborers to take their medication without sacrificing a full day's wage.

However, while 'case holding' (treatment success) exceeded global targets, 'case finding' remained a lingering challenge. During this period, the Case Detection Rate (the estimated percentage of true incident cases diagnosed) hovered around 60%. This indicates that while the system was brilliantly curing those it found, a significant reservoir of TB remained undiagnosed in the community, hampered by a lack of microscopic diagnostic facilities at the extreme periphery.

To fully grasp the magnitude of the DOTS achievement, one must contextualize the socio-economic devastation wrought by Tuberculosis in Nepal. TB is uniquely insidious because it primarily targets the economically active demographic. When a primary breadwinner in an agrarian Nepalese household contracts TB, the economic shockwaves are catastrophic.

Prior to DOTS, the protracted illness, combined with the catastrophic out-of-pocket expenditure associated with seeking care at distant hospitals, frequently drove families into insurmountable debt. Land was sold, children were pulled from school to replace lost agricultural labor, and the household plunged below the poverty line. The 28% default rate observed in 1994 was largely an economic default; patients simply ran out of money to travel to the hospital to collect their next batch of medication.

The DOTS strategy was, therefore, not just a medical intervention; it was a profound anti-poverty program. By decentralizing drug provision, offering the drugs entirely free of charge, and guaranteeing a cure for 87% of patients within 6 to 8 months, the NTP effectively restored the economic engines of thousands of rural households. The reduction in the Treatment Adherence Friction Index (TAFI) directly translated into macroeconomic preservation at the village level.



5. EXPANDED ARCHITECTURAL ANALYSIS

The implementation of the DOTS strategy across Nepal's 75 districts required an unprecedented scale-up of public health infrastructure, specifically in human resources, supply chain management, and health informatics.

A. Human Resources and FCHV Integration

The topography of Nepal dictates that placing a physician or registered nurse in every village is economically and practically impossible. The success of DOTS rested on task-shifting. The National Tuberculosis Programme leveraged the existing network of Female Community Health Volunteers (FCHVs) and Village Health Workers (VHWs). These community members were trained to act as DOTS observers. Because they lived within the same communities as the patients, the distance penalty (D_clinic) in the TAFI equation was functionally reduced to zero. Furthermore, the stigma penalty (F_stigma) was mitigated because patients were receiving care from familiar neighbors rather than institutional authorities.

B. Supply Chain and Logistics (The Push-Pull System)

A critical failure point in pre-1996 TB control was drug stock-outs. If a patient traveled to a hospital and found no medication available, the likelihood of them returning diminished exponentially. To solve this, the DOTS rollout introduced a rigorous 'push-pull' logistics model. District stores operated on a 'pull' system, requesting blister packs from the regional medical stores based on actual consumption data (cohort reporting). However, at the peripheral level, a 'push' system was utilized: for every patient diagnosed and registered, a complete 6-to-8 month course of medication was physically assigned and stored in a designated box with the patient's name on it. This guaranteed that once a patient initiated treatment, their specific drug supply was completely insulated from national stock-out fluctuations.

C. Health Informatics and Cohort Monitoring

The transition from cross-sectional aggregate reporting to longitudinal cohort monitoring was revolutionary. Previously, the HMIS simply counted 'number of patients treated.' The DOTS framework introduced standardized ledgers (the TB Register) that tracked individual patients across time. Every quarter, district supervisors convened to review the outcomes of the cohort registered 12 to 15 months prior. This closed-loop information system allowed managers to identify exactly which Sub-Health Posts were experiencing high default rates and intervene immediately, shifting the entire health system from a reactive to a proactive posture.

D. The Sociotechnical Paradigm

By analyzing the DOTS rollout as a sociotechnical system, it becomes evident that the technology (the multi-drug therapy) was entirely dependent on the social architecture (the FCHVs and the cohort monitoring). The TAFI model proves mathematically that investing in the social architecture yields higher returns on treatment success than investing in marginally better pharmaceuticals without a delivery mechanism.

6. CONCLUSIONS

Nepal's transition from a 40% to an 87% TB treatment success rate within seven years stands as a monumental achievement in health systems engineering. The DOTS strategy proved that organizational friction, not patient ignorance, is the primary driver of treatment default.

To build upon this success and push toward the ultimate elimination of TB, policymakers must adopt the following directives:

A. Expand Microscopic Frontiers: High treatment success is irrelevant for patients who are never diagnosed. Investment must pivot toward expanding quality-assured sputum smear microscopy centers deeper into the rural periphery to improve the Case Detection Rate.

B. Maintain Vigilance Against MDR-TB: While the 4.5% default rate is excellent, it is not zero. The NTP must rapidly integrate specialized DOTS-Plus strategies to identify and manage the emerging threat of Multi-Drug Resistant strains, which require highly complex, 24-month regimens.

C. Mobilize the Private Sector: A significant portion of urban patients initially seek care in the unregulated private sector, where chaotic prescribing practices risk generating drug resistance. The NTP must implement Public-Private Mix (PPM) models to bring private practitioners under the strict umbrella of DOTS protocols.

The TAFI model is not exclusive to Tuberculosis. It can be applied directly to modern infectious disease challenges, such as HIV/AIDS antiretroviral adherence or even contact-tracing compliance during acute respiratory outbreaks like COVID-19. Engineers and public health officials must continuously aim to minimize the numerator (e.g., through telehealth) while maximizing the

denominator (through social safety nets and community support).

7. ACKNOWLEDGEMENTS

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Author contributions

Deependra Paudel: Conceptualization; Data curation; Formal analysis; Investigation; Methodology; Software (Spatial Simulation); Writing original draft; and Writing - review & editing.

The author declares that he has no conflict of interest.

Data Availability

The longitudinal cohort outcomes utilized in this study are derived from the historical Annual Reports of the Department of Health Services (DoHS) and the National Tuberculosis Centre (NTC) of Nepal.

References:

Department of Health Services (DoHS), Ministry of Health, His Majesty's Government of Nepal. (1995-2002). Annual Reports (2051/52 to 2057/58 BS). Kathmandu, Nepal.

National Tuberculosis Centre (NTC). (2001). Annual Report of the National Tuberculosis Programme, Nepal. Thimi, Bhaktapur.

World Health Organization (WHO). (1999). Global tuberculosis control: WHO report 1999. Geneva.

Bam, D. S., et al. (2001). DOTS in Nepal: a success story. The International Journal of Tuberculosis and Lung Disease, 5(11), 1083-1087.

Volmink, J., & Garner, P. (2007). Directly observed therapy for treating tuberculosis. Cochrane Database of Systematic Reviews, (4).

Dye, C., et al. (1999). Consensus statement. Global burden of tuberculosis: estimated incidence, prevalence, and mortality by country. WHO Global Surveillance and Monitoring Project. JAMA, 282(7), 677-686.

Farmer, P. (1997). Social scientists and the new tuberculosis. Social Science & Medicine, 44(3), 347-358.